

OBSERVATIONS ON THE MORPHOLOGY OF *PORPHYRA VIETNAMENSIS* TANAKA & P.H. HO (BANGIALES, RHODOPHYTA) AT VISAKHAPATNAM COAST, INDIA

S. GHOSH AND J. P. KESHRI*

CAS in Botany, The University of Burdwan, Golapbag, Burdwan 713 104

*e-mail : jpkeshri @ indiatimes.com

ABSTRACT

Porphyra vietnamensis Tanaka & P.H. Ho was collected from the east coast (various parts of Visakhapatnam) of India during a low tide status. Detailed morphology of the taxon was studied. The present specimens are smaller than the type material originally described but slightly bigger than the material described from Visakhapatnam coast.

Keywords : *Porphyra vietnamensis*, morphology, Visakhapatnam coast, India.

INTRODUCTION

Porphyra vietnamensis Tanaka & P.H. Ho (Bangiales, Rhodophyta) is an edible red alga (Subha Rao & al. 2007) first described from Viet-Nam (Tanaka & Pham-Hoang Ho, 1962). The particular species of interest is not so frequently found [Viet-Nam (Tanaka & Ho 1962, Ho 1969, Tsutsui & al. 2005)]; Hawaiian Islands (Abbott 1999, McDermid & al. 2005); China (Tseng, 1984); Pakistan (Silva & al.1996)]. Bt rgesen (1937) reported *P. vietnamensis* as *P. tenera* Kjellman from harbour area of Madras. Sreeramulu (1952) defined the plants as *P. naidum* Anderson from Visakhapatnam (Waltair) coast. Ultimately it was confirmed as *P. vietnamensis* by Unameheswara Rao & Sreeramulu (1963). They collected the same specimens from the Visakhapatnam coast in the year 1964 and 1970. Subsequently various other workers also reported this taxon from the west coast, especially from the coastal area of Goa, [Dhargalkar & al. (1981), Sahoo & al. (2006)]. According to Sahoo & al. (2001) *Porphyra vietnamensis* is the most abundant among the all other *Porphyra* species reported from India. During an algal excursion to Visakhapatnam coast the authors recorded the above mentioned taxon and studied its detailed morphology.

MATERIALS AND METHODS

Algal specimens were collected from intertidal zone of various localities of Visakhapatnam coast (Collection No.V-14, V-33, V-38, and V-105) during a low tide state. Field photographs with proper measurements were made by Nikon SLR camera. The collected samples were preserved in 5% formalin solution made with marine water. The collected materials were kept in plastic bags and also in hard plastic transparent containers with detailed field notes. The thalli were observed under a stereo binocular microscope. Digital photographs by Leica DMLB system were made from preserved specimens. Identifications were made following Desikachary & al. (1990). The collected preserved materials along with the slide preparations are stored in Phycology laboratory, The University of Burdwan for future study and reference purpose.

RESULTS AND DISCUSSION

Porphyra vietnamensis Tanaka & P.H. Ho.

(Tanaka & Ho 1962, p.34, f.10-11; Desikachary & al. 1990, p.36, p1.2, f.4)

The plants are membranous, attached to the rocky substratum with the help of slightly circular disc shaped holdfast like attachment organ. The specimens are mostly found along with some other algae like *Enteromorpha flexuosa*, *E. intestinalis*, and *Chaetomorpha aerea*. Sometimes tiny filaments of *Lyngbya cinerescens* were found lodged on the marginal part of the specimen.




Fig. 1-6. *Porphyra vietnamensis* Tanaka & P.H. Ho: Thallus morphology: 1. Mature thallus
 2. Thallus margine showing spines; 3. Enlarged view of marginal spines;
 4. Pear shaped cells at the basal part of the thallus and 5 & 6. Part of cystocarpic thallus in surface view.

The plant species of interest is monostromatic, 1.8-3.2 cm in length and 0.7-1.5 cm broad: monoecious, prominent male gametangial portions along with zygotosporangial patches appear on the same thallus: reproductive patches are observed along the marginal sides. Vegetative and reproductive portions are of almost same thickness; margin wavy and serrated with spinous projections, The tip portions of the spinous projections are slightly pointed to obtuse rounded in nature; vegetative cells in surface view 9.2-11.3 µm in length and 4.5-8.5 µm in breadth; there is a mucilage layer which is 4-5.2µm thick; each cell is slightly globular to angular in shape, with a single chromatophore, somewhat stellate and covers the whole cell partly; carpogonia are of pinkish patches found in marginal as well as apical part and spermatangia showing its identity as slightly pale patched region at the marginal side; here 64 spermatia arranged in four tiers of four each in a spermatangium and the number of carpospores within a carposporangium is eight; the thallus expanding above into soft blade of varying degrees with carpogonium simple with long trichogyne formed by vegetative cells; carpogonium 19.5-33.7 µm in length and 9.6-11.63 µm in breadth, trichogyne 0.4-1.1 µm in thickness, carpogonia are found on marginal to slight inward portions of the thallus.

After comparing with descriptions provided by Tanaka & Ho (1962) for the type species it is concluded that the observed specimens are smaller than the type material but slightly bigger than the specimens described by Umamaheswara Rao & Sreeramulu (1964) from Vishakhapatnam coast.

ACKNOWLEDGEMENTS

The authors are thankful to the Head of the Department of Botany, The University of Burdwan for providing laboratory facilities, to G. Mustafa and Prof. Pranjit Sarma for their support and encouragement in various stages of this work and to Goutam Bala and C. Prakasam for their constant help in various occasions.

REFERENCES

- Aaai-r, I. A. 1999. Marine red algae of the Hawaiian Islands. Honolulu, Hawaii: Bishop Museum Press.
- BORGESSEN, F. 1937. Contribution to a South Indian Marine flora-II. *J. Indian Bot. Soc.* 16: 1-56.
- DESIKACHARY, T. V., V. KRISHNAMURTHY AND M. S. BALAKRISHNAN. 1990. Rhodophyta 11. Taxonomy Part. Part IIA. Madras Science Foundation: Madras, India. 279.
- DHARGALKAR, V. K., V. V. AGADHI AND A. G. UNTAWALE. 1981. Occurrence of *Porphyra vietnamensis* (Bangiales, Rhodophyta) along Goa Coast. *Mali asagar-Bulletin of National Institute of Oceanography India* 14: 75-77.
- MCDERMID, K. J., B. STUCKCKE, AND O. J. HALEAKALA. 2005. Total dietary fiber content in Hawaiian marine algae. *Botanica Marina* 48:437-440.
- Ho, PRAM-HoANG. 1969. Rongbien Vietnam. Marine algae of South Vietnam. Saigon.
- SAHOO, D., P. BAWHA AND N. KUSHWAH. 2006. Developmental studies in *Porphyra vietnamensis*; a high temperature resistant species from the Indian coast. *J. Appl. Phycol.* 18 (3-5): 279-286.
- SAHOO, D. B., NIVEDITA AND DEBASIS. 2006. Seaweeds of Indian Coasts. APH publishing corporation, New Delhi. India. 283.
- SILVA, P. C., P. W. BASSON AND R. L. MOE. 1996. Catalogue of the benthic marine algae of the Indian Ocean. *Univ. of California Publ. Bot.* 79: 1-1259.
- SUBHA RAO, P. V., A. MANTRI VAIBHAB AND K. GANFSAN. 2007. Mineral composition of edible seaweed *Porphyra vietnamensis*. *Food Chemistry* 102 (1): 215-218.
- SREERAMULU, T. 1952. On a *Porphyra* from Waltair Coast. *Sci. and Cult.* 18: 285-286.
- TANAKA, T. AND P. H. Ho. 1962. Notes on some marine algae from Viet-Nam I. *Mem. Fac. Fish., Kagashima Univ.* 11: 24-30.
- TSING, C. K. 1984. Common seaweeds of China. Beijing: Science Press. 1-318.

- TsUTSUI, I., Q. N. HUYBH, H. D. NGUYEN., S. ARAI AND T. YASHIDA. 2005. The common marine plants of the Southern Vietnam. Japan Seaweed Association. 1-250.
- UMAMAHESERWARA, RAO. M. AND T. SREERAMULU. 1963. Vertical Zonation and seasonal variation in the growth of *Porphyra* on Visakhapatnam Coast. *Curr. Science*. 32: 173-174.
- UMAMAHESHWARA RAO, M., AND T. SREERAMULU. 1964. An ecological study of some intertidal algae of the Vishakhapatnam Coast. *J. Ecol.* 52: 595-616.
- UMAMAHESWARA RAO, M AND T.SREERAMULU. 1970. An annotated list of the Marine algae of Visakhapatnam (India). *Bot. J. Linn. Soc.* 63: 23-45.